

Red Thread Disease

Laetisaria fuciformis

A disease of lawns in the late summer and autumn months, especially in lawns of low fertility, especially Nitrogen



Identification

This disease is now thought to be a combination or complex of two fungal species:

- a) *L.fuciformis* - Red Thread - where the red needles are present
- b) *A.fuciformis* - Pink Patch - where the pink mycelium is present

Red Thread is a very common turf disease. It occurs during mid to late summer and autumn and can persist into mild winters, especially if the weather is humid and damp. As with any other lawn disease, early and correct diagnosis is important.

It is commonly seen as little red needles emerging from the leaf of the affected grasses or as a pink 'cotton wool' on the surface of the lawn. Affected areas can be the size of a golf ball but can coalesce into larger areas.

Ill-defined patches of bleached grass. Pink mycelia are often seen visible in morning dew cover. Red needles are present attached to the leaf blades. Needles (stroma) become brittle on drying and serve to spread disease into new areas.

Where has it come from?

Red Thread naturally occurs as a Pathogen within the turf grass seed and ultimately the mature grasses. The grass seed breeders work hard to cultivate species and varieties of grasses that have 'freedom of Red Thread' pathogens.

Buying certified and quality grass seed with known Species and Cultivars in the mixture is a good start! It can occur mostly in turf and lawn areas of low fertility levels, particularly Nitrogen. As grass growth slows down due to a lack of Nitrogen, the disease becomes more prevalent.

Will it kill the lawn, left untreated?

Red Thread does not actually kill the grasses, merely making them look a bit straw like and unsightly, especially once the disease has caused the damage and dried up. Grass plants are rarely totally killed.

Grass species involved

Red Thread Disease can affect most turf grass species, but particularly Red Fescue (*Festuca rubra*) and Perennial rye grass (*Lolium perenne*) although most cool season grasses as found in the United Kingdom can suffer an outbreak.

What are my options on control of the disease?

Improve fertility levels immediately, especially Nitrogen and Potassium. Feeding a lawn little and infrequently is the worst culprit! Adopt a regular planned feeding programme. Summer control can almost always be assisted by the application of Nitrogen. Choose disease resistant cultivars for initial sowings, repair and over sowing. Maintain a soil pH of 6.5-7.0. Ensure the lawn is mown with a mower with a sharp blade so that the grass leaf is not damaged and weakened.

Bayer Garden Lawn Disease Control controls fungal diseases on established lawns, protects against common lawn diseases such as Fusarium patch and Red Thread. It prevents and cures diseases